

How does prognosis change after colorectal cancer surgery?

Dynamic prediction over time

Mohammad Asghari-Jafarabadi, Simon Wilkins, Katherine Goodall, Veena Suresh, Rathin Gosavi, Paul J. McMurrick

Cabrini Research | Monash University | Cabrini Monash University Department of Surgery

REAL DATA.
DYNAMIC INSIGHTS.
BETTER LIVES.

Background

Prognosis after colorectal cancer surgery is not fixed at the time of surgery. It changes as patients remain recurrence-free, develop recurrence, or survive longer.

Methods

We analysed 3,271 patients from the Cabrini Monash Colorectal Neoplasia Database using a dynamic multi-state / landmark framework. Risks of recurrence/metastasis, cancer-related death and non-cancer death were updated at clinically relevant times among patients still alive and recurrence-free. Conventional and machine-learning models were compared on a separate test sample.

Cohort outcomes


Recurrence/
metastasis
479

Cancer-related
deaths
497

Non-cancer
deaths
198

Deaths after
documented
recurrence
302

Postoperative clinical-state occupancy



Outcome-specific drivers

Hazard ratio (95% CI)
(log scale)

Recurrence/
metastasis

Tumour stage (III vs I-II)
Lymph-node ratio (high vs low)
Tumour size (per 1 cm)
Poor differentiation
Age (per 10 years)




Cancer-related
death

Tumour stage (III vs I-II)
Nodal disease (positive vs negative)
Poor differentiation
Lymphovascular invasion
Age (per 10 years)




Non-cancer
death

Age (per 10 years)
ASA physical status (III-IV vs I-II)
Comorbidity (Charlson ≥ 2 vs 0-1)
Male sex (vs female)
Diabetes (vs no diabetes)



Dynamic prediction

A patient's risk profile is updated at each time point and can be very different from the original estimate at surgery.



Key results

- At surgery, 5-year prediction was strongest for cancer-related mortality and more modest for recurrence.
- Patients remaining recurrence-free for 1 or more years had updated outcome risks.
- Cancer-related mortality was driven mainly by tumour stage and nodal disease.
- Non-cancer mortality was driven mainly by age, ASA status and comorbidity.

Conclusion

Postoperative colorectal cancer prognosis should be viewed as a changing clinical trajectory rather than a single baseline estimate. Dynamic prediction can support more personalised surveillance, survivorship care, patient counselling and multidisciplinary follow-up. External validation is required before clinical implementation.



References

- van Houwelingen HC, Putter H. Dynamic Prediction in Clinical Survival Analysis. CRC Press; 2012.
- Putter H, Fiocco M, Geskus RB. Competing risks and multi-state models. Stat Med. 2007;26:2389-2430.

Abbreviations

ASA = American Society of Anesthesiologists physical status