

Feasibility of Live Remote Exercise for Adults with Advanced Cancer and Cachexia

Kelcey A. Bland¹, Mark Trevaskis², Evelyn B. Parr², Ashley Bigaran³, Meinir Krishnasamy^{4,5,6}, Peter Martin^{7,8}, Natasha Michael⁹, Melissa Moore¹⁰, Isaac Rankin¹, Jennifer Weil¹⁰, Luc J.C. van Loon¹¹, Prue Cormie¹², Eva M. Zopf^{2,13}

¹University of British Columbia, Canada, ²Australian Catholic University, Australia, ³Austin Health, Australia, ⁴Peter MacCallum Cancer Centre, Australia, ⁵University of Melbourne, Australia, ⁶Victorian Comprehensive Cancer Centre Alliance, Australia, ⁷Deakin University, Australia, ⁸Barwon Health, Australia, ⁹Notre Dame, Australia, ¹⁰St. Vincent's Hospital, Australia, ¹¹Maastricht University, Netherlands, ¹²Flinders University, Australia, ¹³Cabrini Health, Australia

BACKGROUND

- Cancer cachexia is a multifactorial syndrome characterized by an ongoing loss of skeletal muscle mass (with or without loss of fat mass)¹.
- Cancer cachexia affects 50-80% of people with advanced cancer² and has a profound impact on physical function, symptom burden and quality of life (QoL).
- Effective supportive care interventions remain limited, and the role exercise is mostly unknown, as few exercise trials have targeted adults with advanced cancer and cachexia.

Objective: To assess the **feasibility, acceptability, and safety** of a live remote, tailored exercise intervention for adults with advanced cancer and cachexia, and to explore preliminary effects on physical function and patient-reported outcomes.

METHODS

Adults with metastatic or locally advanced cancer meeting international cachexia criteria received an 8-week, live remote, tailored aerobic and resistance exercise program (3 sessions/week, 30-45 minutes) plus post-exercise protein supplementation (EX) or usual care (UC) (Fig 1).




Figure 1. ACE trial study design

Primary outcomes: Feasibility (recruitment, attrition, adherence, safety) and acceptability. Secondary outcomes: Physical fitness and function (e.g., 30-s sit-to-stand), body composition, physical activity, nutritional status, QoL, fatigue, and psychological outcomes.

RESULTS

Of 107 screened, 33 participants (17 males and 16 females aged 63 ± 12 years) were randomized (EX: n=17, UC: n=16), with 30 completing the 8-week study. The most common cancer type was gastrointestinal (46%) followed by lung (12%) and breast (12%). On average, participant's six-month weight loss was $8.0 \pm 3.1\%$.

Feasibility and safety: All feasibility targets were met (Fig 2).




Figure 2. Feasibility and acceptability outcomes

Acceptability: 100% felt the program met their expectations, was tailored to needs, and was beneficial

Secondary outcomes: No between-group differences. However, clinically meaningful improvements in the EX group but not the UC group over time for :

Physical function (30-s sit-to-stand):
 $\Delta 3.1$ reps, (95% CI 0.4–5.8)

Health-related QoL (SF-36):
 $\Delta 13.3$, (95% CI: 5.6 to 20.9)

Bodily pain (SF-36):
 $\Delta 16.7$, (95% CI: 2.2 to 31.1)

"It gave me a sense of achievement and a sense of purpose"

CONCLUSION

A live remote, tailored exercise intervention was feasible, acceptable, and safe for adults with advanced cancer cachexia

ACKNOWLEDGEMENTS

This work was supported by funding from the Cabrin Foundation

