

Colorectal cancer during pregnancy: a systematic review and pooled case-level analysis

Katherine Goodall¹, Arman Erkan², Mohammad Asghari-Jafarabadi^{1,3,4}, Raymond Yap¹

¹ Cabrini Monash University Department of Surgery, Cabrini Hospital, Malvern ² Northwell Health Physician Partners Center for Advanced IBD Care, Great Neck, NY, USA
³ Dept of Psychiatry, School of Clinical Sciences, Monash University ⁴ Dept of Biochemistry & Molecular Biology, Monash University

PROSPERO CRD420261504748 · No funding · No conflicts of interest

Background

- CRC is the 3rd most diagnosed cancer and 2nd leading cause of cancer death worldwide¹
- CRC in pregnancy is rare; evidence is mostly case reports and small series^{2–5}
- Rising early-onset CRC and maternal age will increase its incidence⁶
- Symptoms mimic normal pregnancy, delaying diagnosis; care must balance mother and foetus
- No CRC-specific evidence-based guidelines exist; prior reviews pooled small cohorts^{7–8}


Aim

To collate the literature and explore maternal and foetal outcomes of CRC diagnosed during pregnancy or the delivery admission.

Methods

- 5 databases + citation search; no date limit
- English, Japanese & Turkish; 2 of 3 reviewers
- Case reports/series (270 records) and cohort studies (49 records) analysed separately
- Logistic regression adjusted for gestational age & stage^{9–10}

PRISMA 2020 flow diagram



Created in BioRender. Goodall, K. (2026) <https://BioRender.com/yi57kgh>

Results



200
publications
(1940–2026)



322
case
records



81.6%
stage III–IV
(case-based)




39.1%
maternal death
(case-based)

Case-based: rectal/rectosigmoid 45.6% · resection 79.5% · systemic therapy 76.3% · 2nd/3rd trimester diagnosis 87.2%

Created in BioRender. Goodall, K. (2026) <https://BioRender.com/et79nhs>

Adjusted associations (exploratory, not causal)



Conclusions

- Advanced stage and late-gestation diagnosis mark the highest maternal risk, yet foetal outcomes are mostly good
- Systemic therapy was not associated with adverse foetal outcomes, consistent with its accepted use from the 2nd trimester
- A high index of suspicion and early MDT input are essential
- Largest case-level synthesis to date; findings are hypothesis-generating



Scan for the full reference list